

Traumatic Dislocation of the Hip in a High School Football Player

Charlotte Yates, William D Bandy, R Dale Blasier

C Yates, PT, PhD, PCS, is Assistant Professor, University of Central Arkansas, and Center for Translational Neuroscience, University of Arkansas Medical School, Little Rock, Ark.

WD Bandy, PT, PhD, SCS, ATC, is Professor, Department of Physical Therapy, University of Central Arkansas, 300 Donaghey Ave, Conway, AR 72035 (USA), and Physical Therapist, Sportsmedicine Plus/Adolescent Center, Arkansas Children's Hospital, Little Rock, Ark. Address all correspondence to Dr Bandy at: billb@uca.edu.

RD Blasier, MD, is Professor of Orthopedic Surgery, Arkansas Children's Hospital and University of Arkansas Medical School, Little Rock, Ark.

[Yates C, Bandy WD, Blasier RD. Traumatic dislocation of the hip in a high school football player. *Phys Ther.* 2008;88:780-788.]

© 2008 American Physical Therapy Association

Background. Although traumatic dislocation of the hip often occurs as a result of automobile accidents, dislocations have been reported to occur during sports activities.

Objective. Using the experience in treating a 17-year-old high school football player with a posterior dislocation, complicated by involvement of the sciatic nerve, this case report provides background information on hip dislocations and provides a description of the immediate treatment by the physician, followed by 6 weeks of immobilization, and a detailed account of the 5-month intervention.

Case Description. The patient was injured while making a tackle during a high school football game when another player fell on him from behind. The case report describes his plan of care after immediate hip reduction surgery and 6 weeks on crutches. Generally, the program utilized a progression of non-weight-bearing resistance training and stretching in the initial stages of intervention and progressed to weight-bearing activities (on land and in the pool) as the patient was able to tolerate more stress. In addition, the treatment of the sciatic nerve using electrical stimulation during treadmill walking is described.

Outcomes. The patient was seen in an outpatient physical therapy clinic an average of 2 times per week for 5 months. At the end of 5 months, results of the Lower Extremity Functional Scale (LEFS) indicated that recreational and sporting activities were within normal limits, and the patient was able to return to playing on his high school football team the next year.



Post a Rapid Response or
find The Bottom Line:
www.ptjournal.org

explanation of injury

Traumatic dislocation of the hip is an orthopedic emergency requiring early recognition and prompt reduction for successful management.^{1,2} In a posterior dislocation, the head of the femur lies posterior to the acetabulum and the injured lower extremity has a clinical presentation of shortening, medial (internal) rotation, flexion, and adduction. In an anterior dislocation, the femoral head lies anterior to the acetabulum and the injured lower extremity has a clinical presentation of abduction and lateral (external) rotation of the hip.³ According to a retrospective review of 62 patients (mean age=34.5 years, range=14–72) with traumatic dislocation of the hip, Sahin et al² reported that 57 (92%) were posterior dislocations and 5 (8%) were anterior dislocations. Posterior dislocation as the most frequent direction of dislocation is well supported in the literature.^{1,3–6}

The vast majority of dislocations occur as a result of automobile accidents.⁵ In their review of hip dislocations in 62 patients, Sahin et al² reported 52 (83.9%) of the patients sustained their hip dislocation due to traffic accidents. The most common mechanism of injury for a hip dislocation during an automobile accident is when the person's knee (with hip flexed) strikes the dashboard, forcing the head of the femur posteriorly over the rim of the acetabulum.⁴ However, if the thigh is abducted, impact on the knee would cause further abduction and lateral rotation of the hip, leading to an anterior dislocation, which occurs less frequently than a posterior dislocation.⁴

rare injury to occur in sports

Traumatic dislocation of the hip rarely occurs in sports activities. Sahin et al² reported that 2 (3.2%) of the hip dislocations in their study were the result of athletics. Lamke⁷ investigated 110 traumatic disloca-

tions of the hip and reported that 5.5% occurred during sports activities. More recently, Chudik et al⁸ estimated that only 2% to 5% of all hip dislocations occurred during participation in sports. The injury tends to occur secondary to a collision in sports such as skiing or football.^{5,9,10} A frequent mechanism of injury for a posterior hip dislocation is the knee striking the ground with the hip in a flexed position, thereby forcing the femoral head posteriorly over the rim of the acetabulum.^{5,9,11}

A review of the literature related to the treatment of patients with traumatic dislocation of the hip reveals frequent discussions of immediate intervention (open versus closed reduction) and the need for frequent follow-up examinations to rule out postinjury complications.^{3–5,9,12} Information and details concerning the appropriate plan of care following immediate intervention are lacking. Regardless of whether the hip dislocation is the result of an automobile accident or participation in athletics, no descriptions of the specific intervention and plan of care exist. Palotta and Andrish, for example, suggested that “return to activity is permitted only when strength, motion, and agility have been achieved.”^{3(p610)} Anderson et al suggested that “a return to sport is allowed if the MRI [magnetic resonance image] is negative and there is pain-free range of motion.”^{5(p526)} The authors of these articles provided no description of the appropriate plan of care after the immediate reduction of the dislocated hip.

Although previous literature exists as to the immediate emergency treatment for an individual with a traumatic dislocation of the hip, the specific plan of care for the rehabilitation of this type of injury is not described. Therefore, the purpose of this case report is to describe the physical therapy plan of care for a

17-year-old high school football player with a posterior hip dislocation complicated by involvement of the sciatic nerve.

Patient History

The patient was a 17-year-old male football player who was injured during a high school football game while making a tackle. The patient later reported that he felt a “pop” and immediate pain in his right hip when making the tackle and turning when another player fell on the back of his thigh. The patient was evaluated on the field and transported to the emergency department via ambulance.

Emergency Department Examination/Intervention

The patient arrived at the emergency department with his right lower extremity propped on pillows in flexion, adduction, and medial rotation—consistent with a posterior dislocation of the hip. In addition, prior to radiographic evaluation and reduction, the patient had decreased light touch sensation in his foot and was unable to flex or extend his toes or ankle, consistent with a sciatic nerve injury. Radiologic evaluation confirmed that he had a right posterior hip dislocation (Fig. 1).

The hip was reduced with the patient in the supine position. After conscious sedation allowed the patient and his musculature to relax, the reduction was performed. The physician applied traction to the flexed knee in line with the axis of the thigh with the hip flexed 45 degrees while an assistant stabilized the body and trunk to allow countertraction. After a few moments of traction, a gentle external rotation force was applied, and the hip was felt to reduce. The hip was put through gentle range of motion (ROM) to assess for crepitus, which could suggest the presence of a retained intra-articular fragment of bone, cartilage,

before a PT can release player—player must be tested



Figure 1.
Radiograph of pelvis taken in emergency department prior to hip reduction.



Figure 2.
Radiograph of pelvis taken in emergency department after hip reduction.

or soft tissue. No crepitus was present. Post-reduction radiographs were taken to ensure that the hip was successfully reduced and that nothing, such as interposed joint capsule or cartilage, was blocking a full reduction. Post-reduction radiographs revealed a good reduction, with no evidence of fracture or avascular necrosis (Fig. 2).

At this time, the patient was referred for physical therapy for instruction on touch-down weight bearing on the right lower extremity. Due to difficulty ambulating with crutches and the amount of pain in the hip, he was admitted to the hospital after reduction of his hip dislocation for intravenous pain control and physical therapy for instructions on

crutches. A computed tomography scan of the right hip was ordered and was completed on the morning of the admission. The results showed a normal scan of a good reduction, with no evidence of bony fragments or fracture about the joint. The day after reduction and admission, the patient was walking independently with crutches, and the pain was better controlled. However, he continued to be limited in active dorsiflexion of the ankle, and a resting ankle-foot orthosis was ordered to maintain the ankle in neutral dorsiflexion when he was in bed. The patient was discharged from the hospital with instructions for touch-down weight bearing, with periodic follow-up with the orthopedic fracture clinic.

Initial Physical Therapist Examination

Presenting Complaint

At the time of the initial examination, 6 weeks after injury, the patient arrived for physical therapy ambulating independently with a Trendelenburg gait when weight bearing on the right lower extremity. His chief complaints were an inability to walk and run without hip pain, weakness in the hip, and difficulty in moving his right foot due to "intense pain" in the calf, as well as intermittent pain in the right posterior thigh.

Step 1: before doing anything must have initial examination with patient

Functional Status

Prior to the examination, the patient completed the Lower Extremity Functional Scale (LEFS).¹³ Binkley et al¹³ have reported the LEFS to be reliable, valid, and sensitive to change. In addition, the LEFS has been used to successfully document functional abilities in patients with osteoarthritis of the hip and knee¹⁴ and community-dwelling patients with difficulties in mobility referred for physical therapy.¹⁵ The LEFS is used to qualitatively assess an individual's functional status during 20 specific functional tasks on a scale

Table.Patient's Lower Extremity Functional Scale (LEFS) Scores at Initial Physical Therapy Visit (Pre) and at Discharge (Post)^akeep
record of
activities
for patient
with initial
and
discharge

Activities	Extreme Difficulty or Unable to Perform		Quite a Bit of Difficulty		Moderate Difficulty		A Little Bit of Difficulty		No Difficulty	
	0 Points		1 Point		2 Points		3 Points		4 Points	
	Pre	Post	Pre	Post	Pre	Post	Pre	Post	Pre	Post
Usual work, household or school activities							▲			★
Usual hobbies, recreational and sporting activities	▲									★
Getting into or out of bath									▲	★
Walking between rooms									▲	★
Putting on shoes or socks									▲	★
Squatting									▲	★
Lifting objects from the floor			▲							★
Light activities around the house							▲			★
Heavy activities around the house			▲							★
Getting in or out of the car							▲			★
Walking 2 blocks			▲							★
Walking a mile	▲									★
Ascending/descending a flight of stairs	▲									★
Standing for an hour					▲					★
Sitting for an hour							▲			★
Running on even ground	▲									★
Running on uneven ground	▲									★
Making sharp turns while running fast	▲									★
Hopping	▲									★

^a ▲=score at initial physical therapy visit; ★=score at physical therapy discharge.

from 0 (unable to perform actively) to 4 (no difficulty). The patient's pre-intervention score on the LEFS was 33/80 (Table).

Systems Review

The patient appeared to be a healthy 17-year-old athletic male. He was 175.26 cm (5 ft 9 in) tall, weighed 77.1 kg (170 lb). Heart rate, blood pressure, and respiration were well within normal limits. Other than an antalgic gait when weight bearing on his right lower extremity, the patient demonstrated normal, coordinated movement and motor function. Gross assessment indicated no scars, edema, or any abnormalities with the skin. Screening for any musculoskeletal abnormalities of the upper ex-

trémities and lumbar spine showed full ROM, substantial strength (force-generating capacity), and no pain with any movements. In addition, gross assessment of sensation with light touch indicated decreased sensation ("tingling") in toes 2 through 5 and on the dorsum and plantar aspect of the right foot.

ROM

Examination indicated full passive ROM bilaterally in the patient's hips, knees, and ankles. The patient complained of pain with gentle overpressure into medial and lateral rotation of the right hip. He was apprehensive when performing a posterior glide of the hip joint. In addition, the patient complained of pain with gen-

tle overpressure to dorsiflexion of the right ankle. He had difficulty moving toes 2 through 5 of the right foot, consistent with a sciatic nerve injury. Figure 3 illustrates the relationship of the acetabulum, the head of the femur, and the posterior location of the sciatic nerve and how a posterior dislocation of the femur can cause pressure on the sciatic nerve. Given the mechanism of injury, it is probable that this athlete had a tension type of injury rather than a compression type of injury of the sciatic nerve.

Pain

A 10-point visual analog scale was used to evaluate the patient's pain with the most pain-provoking activi-




Figure 3. Illustration of the relationship of the acetabulum, the head of the femur, and the posterior location of the sciatic nerve.

ties, with 0 representing “no pain” and 10 representing “the worst pain imaginable.” Previous research^{16,17} has shown this method of pain assessment to be both reliable and valid. The visual analog scale scores were 8/10 with palpation of right calf, 8/10 with isometric hip adduction, and 5/10 with ambulation. Attempting to assess pain during running or hopping activities was not appropriate at the initial examination.

Muscle Strength

Muscle strength was assessed manually using standardized methods and test positions described by Reese.¹⁸ Good reliability and validity can be achieved when using standardized methods, especially when muscle grades are below the level of 4.¹⁸ The patient had excellent strength in his left (uninvolved) lower extremity. Manual muscle testing of the right lower extremity indicated minor weakness during hip flexion (grade 4/5) and knee extension (grade 4/5). More significant weakness was found during the following move-

ments: hip extension (grade 2/5), hip abduction (grade 3/5), hip medial rotation (grade 3/5), hip lateral rotation (grade 3/5), knee flexion (grade 3/5), dorsiflexion (grade 2/5), and plantar flexion (grade 2/5). Hip adduction was not tested due to pain.

Assessment

The patient exhibited right hip pain, decreased muscle strength in the right lower extremity, and poor functional abilities secondary to right posterior hip dislocation. In addition, he appeared to have signs of sciatic nerve complications, as indicated by pain in the posterior thigh and calf, and difficulty flexing and extending his toes, and he reported a tingling sensation with light touch to his toes and the dorsum and plantar aspect of the foot. The goal was to return the patient to the highest level of function possible. The prognosis for return to football was guarded due to the traumatic nature of the injury and the involvement of the sciatic nerve.

Initial Intervention

The initial goals of treatment were to increase the strength of the involved extremity and to improve the quality of gait. A daily home exercise program of isometric adduction, straight leg raises, and gravity-resisted hamstring curls was initiated. The patient was instructed to perform each of these exercises 15 times, 2 times per day. He was instructed to perform an isometric contraction of the hip extensors after performing the hamstring curls, as he was unable to perform hip extension exercises in a prone position without significant recruitment of the paraspinal muscles.

The patient was seen 7 times during the first month. Physical therapy twice weekly focused on: (1) proprioceptive neuromuscular facilitation D1 and D2 patterns¹⁹ (15 repetitions of each pattern); (2) gentle manual stretching of the hamstring muscle in the 90-90 position (5 repetitions for 30 seconds) and ankle dorsiflexion with knee extended (5 repetitions for 30-second holds^{20,21}); (3) proximal strengthening during dynamic activities (5 minutes with rest breaks) in the half-kneel position with the left leg leading; (4) stationary bike training for endurance (15 minutes); (5) progressive resisted exercise of the hip abductor, internal rotators, and external rotators, with and without elastic bands, beginning with no weight and progressing to 2.27 kg (5 lb) at 3 months (10 repetitions × 2 sets); (6) non-weight-bearing strengthening of the plantar flexors²² using an elastic band, beginning with yellow Thera-Band* and progressing to green Thera-Band at 2 months (15 repetitions × 2 sets); and (7) pre-gait activities (weight shift laterally, weight shift in a diagonal pattern with focused attention of plantar-flexor activation at heel-off

provide exercises for patient

focus on different sets per month

* The Hygenic Corp, 1245 Home Ave, Akron, OH 44310-2575.

of right lower extremity during the pre-swing phase, and single-leg stance on the right with the left foot elevated on a secondary surface).

Month 2

Examination. Observation at the beginning of the second month revealed that the patient demonstrated an abnormality at mid-stance and terminal stance due to continued weakness in his plantar flexors, contributing to a decreased step length on the left. He demonstrated an obvious lack of force production during terminal stance, with a decreased heel raise on the right lower extremity compared with the left lower extremity. The patient demonstrated decreased force production of the triceps surae muscle with attempted single-leg heel raises.

Strength improvements from the initial evaluation were: grade 4/5 (from 3/5) for hip medial and lateral rotation and grade 4/5 (from 3/5) for hip abduction. Manual muscle testing revealed muscle strength of grade 3/5 for hip adduction. Circumferential measurements around the gastrocnemius muscle revealed the left calf to be 1½ cm larger than the right calf. The patient demonstrated poor proprioception in the right lower extremity with single-leg stance activities and was only able to maintain his balance for 3 seconds with his eyes closed. The patient's LEFS score at 2 months was 45/80.

Intervention. The patient was seen 7 times during the second month. Physical therapy treatment continued 2 times per week and focused on strengthening of the right lower extremity, functional strengthening activities, gait refinement, proprioception, and cardiovascular endurance. The patient continued his daily home exercise program as previously described. Pool therapy was added one time a week, but success

was limited initially due to his limited water skills prior to injury. Pool therapy focused on prone kickboard activities and deep-water running and jumping. In addition, treatment focused on proprioception work in the pool and progressed to dynamic proprioception work on land. This included single-leg stance on the right lower extremity in chest-deep water with overhead dynamic upper-extremity tasks for 5-minute intervals and progressed to single-leg stance on the right lower extremity with dynamic upper-extremity tasks on land. The patient continued with progressive strengthening activities, including progressive resisted exercise with Thera-Band and proprioceptive neuromuscular facilitation, for plantar flexion and progressed to weight-bearing strengthening^{23,24} of the plantar flexors and hip extensors, with mini-squats and leg-press activities involving lowering from a step (15 repetitions × 3 sets for both activities). Single-leg heel raises with assistance of his arms to decrease compensatory patterns were initiated at this time (10 repetitions × 3 sets).

common treatment used for any injury

To further address the patient's plantar-flexor weakness, electrical stimulation was initiated in combination with treadmill walking at a variety of speeds and inclines.^{25,26} Electrodes were placed on the heads of the gastrocnemius muscles, and a remote switch activated the gastrocnemius muscles at mid-stance and terminal stance, beginning with 5 minutes and progressing to 20 minutes of treadmill activities during each therapy session by the third month. The Empi Respond Select Unit[†] was used with the remote switch, and amplitude was determined by clear visualization of plantar-flexor contraction within limits of patient tolerance prior to initiation

[†] Empi Inc, 599 Cardigan Rd, St Paul, MN 55126-4099.

ation of weight bearing. The plantar flexors were stimulated using a biphasic asymmetrical balanced waveform with a pulse width of 300 microseconds, a pulse rate of 50 pulses per second, and a ramp time of 0.2 second.

treatments have to do with what areas focusing on per month

Month 3

Examination. At the beginning of the third month, the patient was not able to kick across the width of the pool lying prone on a float. Strength testing indicated that all hip muscles had improved to grade 5/5. Circumferential measurements indicated a 1-cm difference between the left and right calves. The patient was able to demonstrate a supine straight leg raise with a 2.27-kg (5-lb) weight for 20 repetitions. His LEFS score at this time improved to 60/80.

Intervention. The patient was seen 7 times during the third month. Given the improvement during the second month, the twice-weekly treatments were progressed to more functional activities, and the frequency of pool therapy was decreased to twice a month. A functional progression program, defined as a series of sport-specific basic movement patterns that are gradually progressed according to the difficulty of the skill and the client's tolerance,²⁷ was initiated. The patient started working on basic plyometric drills consisting of jumping off a 5.08-cm (2-in) mat and stepping back up and progressed to jumping up and down from a 5.08-cm mat for 30 seconds.²⁸

The literature indicates that the "calcaneus everts about 5 degrees and the subtalar joint moves into pronation at loading response."^{29(p14)} As the patient progressed to jogging and running activities on land, we noted that his ankle remained in subtalar neutral to minimal supination, potentially placing him at risk for an inversion ankle sprain. Therefore,

step at the beginning of each month

add more as patient can do more

the electrodes were placed on the lateral leg to activate the peroneal muscles and address the timing of the activation of the peroneal muscles in mid-stance and terminal stance. The patient used the remote switch as he progressively worked on the treadmill at increasing speeds. Two months of focused work corrected the problem.

The final functional activity implemented into the treatment plan was rope jumping (30 repetitions \times 2 sets). The patient continued to address his cardiovascular conditioning on his own through stationary bike training (20–30 minutes, 3 times per week).

Month 4

Examination. At the beginning of the fourth month, the patient had full hip ROM and mild discomfort with maximal overpressures performed at the end range of hip medial rotation and at the end range of hip lateral rotation. His score on the LEFS was 72/80.

The patient also performed functional tests consisting of a vertical jump test and a 3-hop, one-legged cross-over test.²⁷ Functional tests are the performance of one maximal effort of a functional activity, or series of activities, in an attempt to indirectly assess muscle strength and power and to quantify function. The one-legged vertical jump test measured the height that the patient jumped while touching the wall at the height of the jump and landing on the same leg. During the 3-hop, one-legged cross-over test, the patient hopped forward 3 consecutive times while crossing back and forth over a center strip that was 15 cm in width. No differences were found between the right and left limbs for either test, and the patient reported no pain.

Intervention. The goal at this time was to increase the difficulty of the functional activities in preparation for the patient returning to football practice. The patient was instructed in a home exercise program consisting of running 0.4 to 0.8 km (0.25–0.5 mile), jumping rope, running up and down bleachers, 36.6-m (40-yd) sprints (forward, backward, and side-to-side [carioca]), and agility drills in which he performed start-and-stop and cutting activities. He was instructed not to participate in spring football drills or in tackling drills. At this point, the patient was seen weekly to monitor his progress. He was seen 4 times during the fourth month.

Month 5

The patient was seen as part of the pre-participation screening for his football team 2 months prior to the first football practice in the fall. Examination at that time indicated that he was having no difficulty or pain with his exercise program. He still complained of pain with aggressive overpressure at end ranges of medial and lateral rotation in the right hip that was not present in the left hip. His LEFS score was 80/80. At this time, the patient was discharged from the physical therapy service and given permission to fully participate in football in the fall. final step

Follow-up

At follow-up 1 year 3 months after the injury, the patient did not complain of hip pain with any activities. He had participated in all 14 football games of the season, leading to winning a state championship, with no problems and no reinjury. He had no pain with active ROM. He did describe “achy pain” in cold weather. His sciatic nerve function had fully returned. Radiologic examination showed no signs of degenerative changes or avascular necrosis.

Discussion

Traumatic hip dislocation most commonly occurs as a result of an automobile accident in which a person’s knee strikes the dashboard, forcing the head of the femur posteriorly.⁹ The most common mechanism of injury in the rare cases of hip dislocation in the athlete is a forward fall on the knee with a flexed hip, forcing the head of the femur posteriorly over the rim of the acetabulum.⁴ Giza et al¹² suggested a second mechanism of injury for an athlete receiving a hip dislocation during sports activities—a blow from behind when the athlete is on all 4 limbs. This type of injury appears to be what occurred to the patient in this case report. The athlete was making a tackle and received a blow to the thigh from behind.

To date, no other publication has provided a detailed plan of care for an athlete following a posterior dislocation of the hip. Following immediate reduction and 6 weeks on crutches, the program described in this case report utilized non-weight-bearing resistance training and stretching in the initial stages of intervention and progressed to weight-bearing activities as the patient was able to tolerate more stress. The use of pool therapy allowed earlier weight-bearing activities due to the buoyancy of the water. Throughout the intervention, the pain in the hip was closely monitored and, if present, the patient was not progressed to a higher level of work.

This positive outcome does not discount the possibility that our patient continues to be at risk for complications related to his injury, with the most significant concern being the development of degenerative arthritis and avascular necrosis of the femoral head. Degenerative arthritis is thought to be caused or accelerated by scuffing of the smooth joint cartilage during the dislocation reduction

event, and it cannot be completely eliminated or reversed. The most significant concern following a dislocated hip is avascular necrosis of the femoral head. This condition can occur because the important blood vessels that supply the femoral head become torn or stretched during dislocation. The femoral head loses its blood supply, which leads to degeneration, and eventually the joint becomes severely arthritic. No reliable cure exists for this unfortunate condition. Avascular necrosis occurs in 10% to 20% of patients.^{2,3,5}

Time to reduction of the hip also is an important factor to consider in the motor nerve damage associated with nerve injury.³⁰ Our patient was fortunate in the quality of his on-the-field management, the regional children's hospital being within a 10-minute drive of his high school, and the ability of the orthopedic surgeon to reduce the hip in the emergency department. This athlete was at a lower risk for additional complications of labrum damage because bony fragments were not detected on the computed tomography scan at the 1-day follow-up and he denied any pain with hip active ROM.^{2,3,5}

The incidence of nerve injury following a posterior hip dislocation varies. In a review of the literature on traumatic dislocation of the hip, Cornwell and Radomisli reported that the incidence among adults ranged from 0% to 20%, "with the majority reporting incidences in the range of 10%-15%."^{31(p86)} However, a review of the literature by Giannoudi et al³² indicated that only one case of sciatic nerve complication has been documented following an injury due to sports. In that study, Tennent et al³³ described a 22-year-old basketball player who slipped when landing from a jump shot, doing the splits, and sustaining a posterior dislocation of the hip, resulting in sci-

atic nerve palsy. Therefore, the present case report is unique in that the patient sustained a complication to the sciatic nerve.

The mechanism for the injury to the sciatic nerve may be the dislocated femoral head compressing the nerve.³¹ Epstein⁴ suggested that the circumduction motion during the closed reduction of the posteriorly displaced femoral head may produce a traction injury of the sciatic nerve. Cornwell and Radomisli³¹ described the symptoms of a neurologic injury after a posterior hip dislocation to include pain, paresthesia, and weakness in the distribution of the affected nerve. These symptoms are consistent with the clinical presentation of the patient described in the present case report.

Clancy et al³⁴ suggested that the majority of the force for plantar flexion of the ankle is generated by the gastrocnemius-soleus muscle complex and that inappropriate timing of the plantar-flexion contraction during terminal stance may cause inefficient walking. In addition, Marqueste et al³⁵ indicated that treadmill walking or running performed when a muscle is reinnervating causes positive effects on histochemical muscle fiber alterations, contractile properties, enzyme activities, and muscle weight. Furthermore, Marqueste et al³⁵ suggested that chronic electrical stimulation of denervated muscle is known to accelerate the recovery of normal function in reinnervated muscle fibers.

^{add more activities gradually}
As the patient's condition allowed, a series of graduated activities were introduced in order to progressively place more and more demand on the patient's hip. The progressive increase in activity involved an increase in loading on the tissues in an activity-specific fashion. An intimate knowledge of the sport and the specific duties required of the athlete for

playing football were important for a successful functional progression program.

Finally, no elaborate strength measurements were taken. The clinician relied on manual muscle testing (ensuring that all muscles achieved a grade of 5/5) and functional tests. Given that previous research^{36,37} demonstrated that greater sensitivity occurs when 2 one-legged tasks are performed, our patient was required to perform 2 maximal hop tests on the involved (right) leg to the same degree as the uninvolved (left) limb prior to returning to competitive sports.

Conclusion

This case report highlights the plan of care provided to a 17-year-old high school football player following a traumatic dislocation of the hip, complicated by sciatic nerve involvement. Following immediate reduction of the hip by the physician, 6 weeks on crutches, and 5 months of physical therapy intervention, the athlete was able to return to his previous high level of activity and complete a full season as a participant on his high school football team.

All authors provided concept/idea/project design, writing, data collection, project management, and patient. The authors acknowledge Dr Tiffany Huitt for the illustration of the posterior dislocation of the sciatic nerve used in Figure 3.

A platform presentation of this work was given at the Combined Sections Meeting of the American Physical Therapy Association; February 1–5, 2006; San Diego, Calif.

Support was provided to Dr Yates through National Institutes of Health grant RR020146 to the Center for Translational Neuroscience.

This article was received December 19, 2007, and was accepted February 20, 2008.

DOI: 10.2522/ptj.20070298

References

- 1 Yang EC, Cornwell R. Initial treatment of traumatic hip dislocation in the adult. *Clin Orthop Relat Res*. 2000;377:24–31.

- 2 Sahin V, Karakas ES, Aksu S, et al. Traumatic dislocation and fracture-dislocation of the hip: a long-term follow-up study. *Journal of Trauma: Injury Infection and Critical Care*. 2003;54:520-529.
- 3 Paletta GA Jr, Andrish JT. Injuries about the hip and pelvis in the young athlete. *Clin Sports Med*. 1995;14:591-628.
- 4 Epstein HC. Traumatic dislocations of the hip. *Clin Orthop Relat Res*. 1973;92:116-142.
- 5 Anderson K, Strickland SM, Warren R. Hip and groin injuries in athletes. *Am J Sports Med*. 2001;29:521-533.
- 6 Yang RS, Tsuang YH, Hang YS, et al. Traumatic dislocation of the hip. *Clin Orthop Relat Res*. 1991;265:218.
- 7 Lamke L. Traumatic dislocations of the hip. *Acta Orthop Scand*. 1970;41:188-198.
- 8 Chudik S, Answorth A, Lopez V, et al. Hip dislocations in athletes. *Sports Med Arthrosc Rev*. 2002;10:123-133.
- 9 Moorman CT, Warren RF, Hershman EB, et al. Traumatic posterior hip subluxation in American football. *J Bone Joint Surg Am*. 2003;85:1190-1196.
- 10 Matsumoto K, Sumi H, Sumi Y, et al. An analysis of hip dislocations among snow boarders and skiers: a 10-year prospective study from 1992-2002. *J Trauma*. 2003;55:946-948.
- 11 Cooper DE, Warren RF, Barnes R. Traumatic subluxation of the hip resulting in aseptic necrosis and chondrolysis in a professional football player. *Am J Sports Med*. 1991;19:322-324.
- 12 Giza E, Mithöfer K, Matthews H, Vrahas M. Hip fracture-dislocation in football: a report of two cases and review of the literature. *Br J Sports Med*. 2004;38:E17.
- 13 Binkley JM, Stratford PW, Lott SA, et al. The Lower Extremity Functional Scale (LEFS): scale development, measurement properties, and clinical application. *Phys Ther*. 1999;79:371-383.
- 14 Stratford PW, Kennedy DM, Hanna SE. Condition-specific Western Ontario McMaster Osteoarthritis Index was not superior to region-specific Lower Extremity Functional Scale at detecting change. *J Clin Epidemiol*. 2004;57:1025-1032.
- 15 White LJ, Straube D, Keehn MT. Using compensations to assess physical performance for ambulatory patients. *Arch Phys Med Rehabil*. 2004;85:1519-1524.
- 16 Downie WW, Leatham PA, Rhind VM, et al. Studies with pain rating scales. *Ann Rheum Dis*. 1978;37:378-381.
- 17 Price DD, McGrath PA, Rafii A, et al. The validation of visual analogue scales as ratio scale measures for chronic and experimental pain. *Pain*. 1983;17:45-56.
- 18 Reese NB. *Muscle and Sensory Testing*. 2nd ed. St Louis, Mo: Elsevier Saunders; 2005.
- 19 Hanson C. Proprioceptive neuromuscular facilitation. In: Hall CM, Brody LT, eds. *Therapeutic Exercise: Moving Toward Function*. 2nd ed. Baltimore, Md: Lippincott Williams & Wilkins; 2005: chap 16.
- 20 Bandy WD, Irion JM, Briggler M. The effect of time and frequency of static stretching on flexibility of the hamstring muscles. *Phys Ther*. 1997;77:1090-1096.
- 21 Nelson RT, Bandy WD. Eccentric training and static stretching improve hamstring flexibility of high school males. *J Athl Train*. 2004;39:31-35.
- 22 Monahan JP, Hartley R, Hall C, Smith S. The ankle and foot. In: Hall CM, Brody LT, eds. *Therapeutic Exercise: Moving Toward Function*. 2nd ed. Baltimore, Md: Lippincott Williams & Wilkins; 2005: chap 22.
- 23 Davies GJ, Heiderscheit BC, Manske R, et al. The scientific and clinical rationale for the integrated approach to open and closed kinetic chain rehabilitation. *Orthop Phys Ther Clin North Am*. 2000;9:247-267.
- 24 Straker JS, Stuhler PJ. Clinical application of closed kinetic chain exercises in the lower extremities. *Orthop Phys Ther Clin North Am*. 2000;9:185-207.
- 25 Carmick J. Clinical use of neuromuscular electrical stimulation for children with cerebral palsy, part 1: lower extremity. *Phys Ther*. 1993;73:505-513.
- 26 Dubowitz L, Finnie N, Hyde SA, et al. Improvement of muscle performance by chronic stimulation in children with cerebral palsy. *Lancet*. 1988;12:587-588.
- 27 Bandy WB, Rusche KR, Tekulve FR. Reliability and limb symmetry for five unilateral functional tests of the lower extremities. *Isokinet Exerc Sci*. 1994;4:108-111.
- 28 Hall C, Brody LT. Impairment in muscle performance. In: Hall CM, Brody LT, eds. *Therapeutic Exercise: Moving Toward Function*. 2nd ed. Baltimore, Md: Lippincott Williams & Wilkins; 2005: chap 5.
- 29 The Pathokinesiology Service and the Physical Therapy Department of Ranchos Los Amigos National Rehabilitation Center. *Observational Gait Analysis*. Downey, Calif: Los Amigos Research and Education Institute; 2001.
- 30 Hillyard RF, Fox J. Sciatic nerve injuries associated with traumatic posterior hip dislocations. *Am J Emerg Med*. 2003;545-548.
- 31 Cornwall R, Radomisl TE. Nerve injury in traumatic dislocation of the hip. *Clin Orthop Relat Res*. 2000;377:84-91.
- 32 Giannoudi PV, Zelle BA, Kamath RP, Pape HC. Posterior fracture-dislocation of the hip in sports. *Eur J Trauma*. 2003;29:399-402.
- 33 Tennent TD, Chamblor AF, Rossouw DJ. Posterior dislocation of the hip while playing basketball. *Br J Sports Med*. 1998;32:342-343.
- 34 Clancy EA, Cairns KD, Riley PO, et al. Effects of treadmill walking speed on lateral gastrocnemius muscle firing. *Am J Phys Med Rehabil*. 2004;83:507-514.
- 35 Marqueste T, Alliez JR, Alluin O, et al. Neuromuscular rehabilitation by treadmill running or electrical stimulation after peripheral nerve injury and repair. *J Appl Physiol*. 2004;96:1988-1995.
- 36 Tegner Y, Lysholm J, Lysholm M, et al. A performance test to monitor rehabilitation and evaluate anterior cruciate ligament injuries. *Am J Sports Med*. 1986;17:156-159.
- 37 Barber SD, Noves FR, Mangine RE, et al. Quantitative assessment of functional limitations in normal and anterior cruciate ligament-deficient knees. *Clin Orthop Relat Res*. 1990;255:206-214.